

TORSIONLESS MODULES

APOSTOLOS BELIGIANNIS

*Lecture Notes for the Conference
“The Representation Dimension of Artin Algebras”
1-4 May, 2008, Bielefeld, Germany.*

1. THE CATEGORY OF TORSIONLESS MODULES

Let Λ be an Artin algebra and $\mathbf{mod}\text{-}\Lambda$ be the category of finitely generated right Λ -modules. We denote by $\underline{\mathbf{mod}}\text{-}\Lambda$ the stable category of $\mathbf{mod}\text{-}\Lambda$ modulo projectives and by $\overline{\mathbf{mod}}\text{-}\Lambda$ the stable category of $\mathbf{mod}\text{-}\Lambda$ modulo injectives. We denote by \mathcal{P}_Λ , resp. \mathcal{I}_Λ , the category of finitely generated projective, resp. injective, Λ -modules.

Let $\mathbf{Sub}(\Lambda)$ be the full subcategory of $\mathbf{mod}\text{-}\Lambda$ consisting of the submodules of the projectives and let $\mathbf{Fac}(\mathbf{D}\Lambda)$ be the full subcategory of $\mathbf{mod}\text{-}\Lambda$ consisting of the factors of the injectives.

Observation 1: There are adjoint pairs

$$(\Sigma_{\mathbf{P}}, \Omega) : \underline{\mathbf{mod}}\text{-}\Lambda \rightleftarrows \underline{\mathbf{mod}}\text{-}\Lambda \quad \text{and} \quad (\Sigma, \Omega_{\mathbf{I}}) : \overline{\mathbf{mod}}\text{-}\Lambda \rightleftarrows \overline{\mathbf{mod}}\text{-}\Lambda$$

where Ω is the usual syzygy functor and Σ is the usual cosyzygy functor.

Indeed $\Sigma_{\mathbf{P}}(\underline{A})$ and $\Omega_{\mathbf{I}}(\overline{A})$ are defined by the exact sequences:

$$A \xrightarrow{f^A} P_0^A \rightarrow \Sigma_{\mathbf{P}}(A) \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \Omega_{\mathbf{I}}(A) \rightarrow I_A^0 \xrightarrow{f_A} A$$

where f^A is the minimal left \mathcal{P}_Λ -approximation of A and f_A is the minimal right \mathcal{I}_Λ -approximation of A . The above approximations exist since \mathcal{P}_Λ and \mathcal{I}_Λ are of finite type and can be described explicitly.

Indeed an easy computation shows that

$$\Sigma_{\mathbf{P}} = \text{Tr}\Omega\text{Tr} \quad \text{and} \quad \Omega_{\mathbf{I}} = \text{DTr}\Omega\text{TrD}$$

where D is the usual duality.

Observation 2: Clearly $\text{Im}(f^A) = \Omega\Sigma_{\mathbf{P}}(A)$ and $\text{Im}(f_A) = \Sigma\Omega_{\mathbf{I}}(A)$. Moreover:

- (i) $\Omega(\underline{\mathbf{mod}}\text{-}\Lambda)$ is a reflective subcategory of $\underline{\mathbf{mod}}\text{-}\Lambda$ with reflection the map $\underline{A} \rightarrow \Omega\Sigma_{\mathbf{P}}(\underline{A})$. In particular:

\underline{A} lies in $\Omega(\underline{\mathbf{mod}}\text{-}\Lambda)$ iff the map $\underline{A} \rightarrow \Omega\Sigma_{\mathbf{P}}(\underline{A})$ is invertible

Note that we may choose the map $A \rightarrow \Omega\Sigma_{\mathbf{P}}(A)$ to be epic in $\mathbf{mod}\text{-}\Lambda$ and then the map

$$A \twoheadrightarrow \Omega\Sigma_{\mathbf{P}}(A)$$

is a left $\Omega(\underline{\mathbf{mod}}\text{-}\Lambda)$ -approximation of A .

- (ii) $\Sigma(\overline{\mathbf{mod}}\text{-}\Lambda)$ is a coreflective subcategory of $\overline{\mathbf{mod}}\text{-}\Lambda$ with coreflection the map $\Sigma\Omega_{\mathbf{I}}(\overline{A}) \rightarrow \overline{A}$. In particular:

\overline{A} lies in $\Sigma(\overline{\mathbf{mod}}\text{-}\Lambda)$ iff the map $\Sigma\Omega_{\mathbf{I}}(\overline{A}) \rightarrow \overline{A}$ is invertible

Note that we may choose the map $\Sigma\Omega_{\mathbf{I}}(A) \rightarrow A$ to be monic in $\mathbf{mod}\text{-}\Lambda$ and then the map

$$\Sigma\Omega_{\mathbf{I}}(A) \twoheadrightarrow A \quad (1.1)$$

is a right $\Sigma(\mathbf{mod}\text{-}\Lambda)$ -approximation of A .

Observation 3: The functors and $\Omega\text{Tr} : \underline{\mathbf{mod}}\text{-}\Lambda \rightarrow \underline{\mathbf{mod}}\text{-}\Lambda^{\text{op}}$ and $\Omega\text{Tr} : \underline{\mathbf{mod}}\text{-}\Lambda^{\text{op}} \rightarrow \underline{\mathbf{mod}}\text{-}\Lambda^{\text{op}}$ induce a duality:

$$\Omega\text{Tr} : \Omega(\underline{\mathbf{mod}}\text{-}\Lambda)^{\text{op}} \xrightarrow{\cong} \Omega(\underline{\mathbf{mod}}\text{-}\Lambda^{\text{op}})$$

with quasi-inverse the functor $\Omega\text{Tr} : \Omega(\underline{\mathbf{mod}}\text{-}\Lambda^{\text{op}})^{\text{op}} \rightarrow \Omega(\underline{\mathbf{mod}}\text{-}\Lambda)$. Indeed this follows from **Observation 2(i)**:

\underline{A} lies in $\Omega(\underline{\mathbf{mod}}\text{-}\Lambda)$ or $\Omega(\underline{\mathbf{mod}}\text{-}\Lambda^{\text{op}})$ iff the map $\underline{A} \rightarrow \Omega\Sigma_{\mathbf{P}}(\underline{A})$ is invertible iff the map $\underline{A} \rightarrow \Omega\text{Tr}\Omega\text{Tr}(\underline{A})$ is invertible.

The duality ΩTr is the functor η constructed in [1].

Observation 4: Since clearly $\underline{\text{Sub}}(\Lambda) = \Omega(\underline{\mathbf{mod}}\text{-}\Lambda)$ and $\overline{\text{Fac}}(\overline{\text{D}}(\Lambda)) = \Sigma(\overline{\mathbf{mod}}\text{-}\Lambda)$, composing the above duality with $\overline{\text{D}}$, we obtain an equivalence:

$$\overline{\text{D}}\Omega\text{Tr} : \underline{\text{Sub}}(\Lambda) \xrightarrow{\cong} \overline{\text{Fac}}(\overline{\text{D}}(\Lambda))$$

with quasi-inverse the functor $\Omega\text{Tr}\overline{\text{D}}$.

Let A be in $\mathbf{mod}\text{-}\Lambda$ and consider the following exact sequences

$$0 \longrightarrow A \longrightarrow I \longrightarrow \Sigma(A) \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow \Omega(I) \longrightarrow P \longrightarrow I \longrightarrow 0$$

where $A \rightarrow I$ is the injective envelope of A and $P \rightarrow I$ is the projective cover of I . The above exact sequences induce the following exact sequence:

$$0 \longrightarrow \Omega(I) \longrightarrow \Omega\Sigma(A) \longrightarrow A \longrightarrow 0$$

and it is easy to see that the epimorphism

$$\Omega\Sigma(A) \twoheadrightarrow A \quad (1.2)$$

is a right $\Omega(\mathbf{mod}\text{-}\Lambda) = \underline{\text{Sub}}(\Lambda)$ -approximation of A .

Now consider the full subcategory

$$\underline{\text{Sub}}(\Lambda) \oplus \overline{\text{Fac}}(\overline{\text{D}}(\Lambda)) := \text{add} \{ X \oplus Y \in \mathbf{mod}\text{-}\Lambda \mid X \in \underline{\text{Sub}}(\Lambda), Y \in \overline{\text{Fac}}(\overline{\text{D}}(\Lambda)) \}$$

which is a generating and cogenerating subcategory of $\mathbf{mod}\text{-}\Lambda$.

Also consider the pull-back of the maps (1.1) and (1.2):

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega(I) & \longrightarrow & X & \longrightarrow & \Sigma\Omega_{\mathbf{I}}(A) \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \Omega(I) & \longrightarrow & \Omega\Sigma(A) & \longrightarrow & A \longrightarrow 0 \end{array}$$

Then the map $X \rightarrow \Omega\Sigma(A)$ is a monomorphism and there is induced a short exact sequence

$$0 \longrightarrow X \longrightarrow \Omega\Sigma(A) \oplus \Sigma\Omega_{\mathbf{I}}(A) \longrightarrow A \longrightarrow 0 \quad (1.3)$$

By using the above pull-back diagram it is easy to see that the map $\Omega\Sigma(A) \oplus \Sigma\Omega_{\mathbf{I}}(A) \rightarrow A$ is a right $\underline{\text{Sub}}(\Lambda) \oplus \overline{\text{Fac}}(\overline{\text{D}}(\Lambda))$ -approximation of A and $X \in \underline{\text{Sub}}(\Lambda) \subseteq \underline{\text{Sub}}(\Lambda) \oplus \overline{\text{Fac}}(\overline{\text{D}}(\Lambda))$.

It follows that $\underline{\text{Sub}}(\Lambda) \oplus \overline{\text{Fac}}(\overline{\text{D}}(\Lambda))$ is contravariantly finite in $\mathbf{mod}\text{-}\Lambda$. This implies that the category

$$\mathbf{mod}\text{-}[\underline{\text{Sub}}(\Lambda) \oplus \overline{\text{Fac}}(\overline{\text{D}}(\Lambda))]$$

of coherent functors $[\underline{\text{Sub}}(\Lambda) \oplus \overline{\text{Fac}}(\overline{\text{D}}(\Lambda))]^{\text{op}} \rightarrow \mathcal{A}b$ is abelian. Consider the restricted Yoneda functor

$$\mathbf{Y} : \mathbf{mod}\text{-}\Lambda \longrightarrow \mathbf{mod}\text{-}[\underline{\text{Sub}}(\Lambda) \oplus \overline{\text{Fac}}(\overline{\text{D}}(\Lambda))], \quad \mathbf{Y}(A) = \text{Hom}_{\Lambda}(-, A)|_{\underline{\text{Sub}}(\Lambda) \oplus \overline{\text{Fac}}(\overline{\text{D}}(\Lambda))}$$

Clearly any coherent functor $F : [\text{Sub}(\Lambda) \oplus \text{Fac}(\text{D}(\Lambda))]^{\text{op}} \rightarrow \mathcal{A}b$ admits a presentation

$$0 \rightarrow Y(A) \rightarrow Y(X_1) \rightarrow Y(X_0) \rightarrow F \rightarrow 0$$

where the X_i lie in $\text{Sub}(\Lambda) \oplus \text{Fac}(\text{D}(\Lambda))$. On the other hand (1.3) induces an exact sequence

$$0 \rightarrow Y(X) \rightarrow Y(\Omega\Sigma(A) \oplus \Sigma\Omega_{\mathbf{I}}(A)) \rightarrow Y(A) \rightarrow 0$$

which implies that $\text{pd } Y(A) \leq 1$. Putting things together we infer that in general:

$$\text{gl. dim mod-}[\text{Sub}(\Lambda) \oplus \text{Fac}(\text{D}(\Lambda))] \leq 3$$

Of course if $\text{Sub}(\Lambda)$ is of finite type, i.e. $\text{Sub}(\Lambda) = \text{add } X$ for some module X , or equivalently $\text{Fac}(\text{D}(\Lambda))$ is of finite type, i.e. $\text{Fac}(\text{D}(\Lambda)) = \text{add } Y$ for some module Y , where we may choose $Y = \Sigma\text{DTr}X$, then $\text{Sub}(\Lambda) \oplus \text{Fac}(\text{D}(\Lambda)) = \text{add}(X \oplus Y)$, the module $X \oplus Y$ is a generator-cogenerator and $\text{mod-}[\text{Sub}(\Lambda) \oplus \text{Fac}(\text{D}(\Lambda))] \approx \text{mod-End}_{\Lambda}(X \oplus Y)$.

It follows that $\text{gl. dim End}_{\Lambda}(X \oplus Y) \leq 3$ and then: $\text{rep. dim } \Lambda \leq 3$.

2. PROBLEMS

Recall that for an Artin algebra Λ , the full subcategory $\text{CM}(\Lambda)$ of Cohen-Macaulay modules, consists of all modules A such that $A \cong \text{Im}(P^{-1} \rightarrow P^0)$ for an infinite acyclic complex $\cdots \rightarrow P^{-1} \rightarrow P^0 \rightarrow P^1 \rightarrow \cdots$ of finitely generated projectives which remains exact after the application of the functor $\text{Hom}_{\Lambda}(-, \Lambda)$. An Artin algebra Λ is said to be of *finite Cohen-Macaulay type*, if $\text{CM}(\Lambda)$ is of finite type.

Recall that Λ is called *Gorenstein*, if $\text{id } \Lambda_{\Lambda} = \text{id } {}_{\Lambda}\Lambda = d < \infty$. For such algebras we have $\Omega^d(\text{mod-}\Lambda) = \text{CM}(\Lambda)$. If Λ is of finite global dimension d , then Λ is Gorenstein and $\Omega^d(\text{mod-}\Lambda) = \text{CM}(\text{mod-}\Lambda) = \mathcal{P}_{\Lambda}$. If Λ is self-injective, then Λ is Gorenstein and $\Omega^0(\text{mod-}\Lambda) = \text{CM}(\text{mod-}\Lambda) = \text{mod-}\Lambda$.

Remark 2.1. If Λ is a 1-Gorenstein algebra of finite Cohen-Macaulay type, then $\text{rep. dim } \Lambda \leq 3$ since in this case $\text{CM}(\Lambda) = \text{Sub}(\Lambda)$. Now the results of [1] suggest to look at the following problems which generalizes the torsionless finite situation.

- (I) If $\Omega^d(\text{mod-}\Lambda)$, for $d \geq 1$, is of finite type, is $\text{rep. dim } \Lambda \leq d + 2$?
If the answer to (I) is yes, then it would follow that:
 - (i) If $\text{gl. dim } \Lambda = d < \infty$, then $\text{rep. dim } \Lambda \leq \text{gl. dim } \Lambda + 2$.
 - (ii) More generally if Λ is d -Gorenstein of finite Cohen-Macaulay type, then $\text{rep. dim } \Lambda \leq d + 2$.
- (II) If the category of d -torsion-free modules in the sense of Auslander (i.e. modules A admitting an exact sequence $0 \rightarrow A \rightarrow P^0 \rightarrow P^1 \rightarrow \cdots \rightarrow P^d$, where the P^i are projective, such that the sequence $\text{Hom}_{\Lambda}(P^d, \Lambda) \rightarrow \cdots \rightarrow \text{Hom}_{\Lambda}(P^1, \Lambda) \rightarrow \text{Hom}_{\Lambda}(P^0, \Lambda) \rightarrow \text{Hom}_{\Lambda}(A, \Lambda) \rightarrow 0$ is exact), is of finite type, is $\text{rep. dim } \Lambda \leq d + 2$?

REFERENCES

- [1] C.M. RINGEL, *The torsionless modules of an Artin algebra*, preprint, (2008).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOANNINA, 45110 IOANNINA, GREECE
E-mail address: abeligia@cc.uoi.gr